
Spatially resolved similarity mapping pinpoints areas of reliable and distinct information across the cortical surface

Joshua Faskowitz[1], Peter A. Bandettini[1,2], Javier Gonzalez-Castillo[1]

[1]Section on Functional Imaging Methods,
[2] Functional Magnetic Resonance Imaging Core Facility,
National Institute of Mental Health, National Institutes of Health, Bethesda, USA

Introduction




- Brain mapping charts features across the cortex
- It's common practice to compare mapped data to:
  - a canonical reference map
  - data from a different modality[5,7]
  - mapped data from other ses. or subs.[4,8]
- Comparing maps is often performed by taking the ***Pearson correlation***
  - We previously[1] demonstrated how to "unwrap" correlation to render ***edge time series***
  - The correlation unwrapping can be applied for spatial correlations (see viz., left)

Map of point-wise contribution to the overall spatial similarity btwn. input data sources

$\frac{\sum}{\text{num. points}}$ = spatial correlation




Spatial similarity map




large values: magnitudes are high or low together

low values: magnitudes are discordant


- Here, describe the process of rendering ***spatial similarity maps*** and demonstrate how the added spatial information can provide additional insights for brain mapping analyses.



1 Map to map



2 Maps to atlas



3 Within-subj. compare


Scenarios where we demonstrate how spatial similarity maps can enhance our understanding of map comparisons

Methods


- Human Connectome Project[6] quality-controlled subset of 352 subjects
- Resting-state data (TR=0.72, 1200 TRs, ~15 min) minimally pre-processed, motion, distortion, ICA-FIX corrected, MSM-aligned, and projected to fsLR surface; bandpass (0.008 - 0.2 Hz) filtered, mean WM, mean CSF, linear, and quadratic drift removed via linear regression in one step using AfNI's 3dTproject
- REST1_LR, REST1_RL averaged for *day 1* data; REST2_RL and REST2_LR averaged for *day 2*
- Time series in fsLR space interpolated to on-avg[2] space using distance-weighted nearest-neighbors
- Structurally derived MSM-aligned data (processed by HCP) interpolated from fsLR space to on-avg space
- Within on-avg space, vertices near corpus callosum were masked out to remove edge effects in feature maps, making for a total of 18,948 vertices of surface used
- neuromaps[10] downloaded from repository, transformed to fsLR w/ included tools, and interpolated to on-avg space

Results


neuromaps atlas correlations




Binary thresholded




Example pairwise relationships



We downloaded all maps projected to on-avg space (n=82). From this matrix, we identified the highest magnitude set of 5 edges within 0.002 of a each other, and rendered their spatial similarity maps.






- This research was made possible thanks to the support of the NIMH Intramural Research Program ZIA-MH002783.
- This work utilized the computational resources of the NIH HPC Biowulf cluster (hpc.nih.gov).
- Thanks to the Section on Functional Imaging Methods for their support and camaraderie


References

1. Faskowitz, J., Esfahlani, F. Z., Jo, Y., Sporns, O., & Betzel, R. F. (2020). Edge-centric functional network representations of human cerebral cortex reveal overlapping system-level architecture. Nature neuroscience, 23(12), 1644-1654.
2. Feilong, M., Jiahui, G., Gobbini, M. I., & Haxby, J. V. (2024). A cortical surface template for human neuroscience. Nature methods, 21(9), 1736-1742.
3. Finn, E. S., Glerean, E., Khojandi, A. Y., Nielson, D., Molfese, P. J., Handwerker, D. A., & Bandettini, P. A. (2020). Idiosynchrony: From shared responses to individual differences during naturalistic neuroimaging. NeuroImage, 215, 116828.
4. Finn, E. S., Shen, X., Scheinost, D., Rosenberg, M. D., Huang, J., Chun, M. M., ... & Constable, R. T. (2015). Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity. Nature neuroscience, 18(11), 1664-1671.
5. Glasser, M. F., & Van Essen, D. C. (2011). Mapping human cortical areas in vivo based on myelin content as revealed by T1-and T2-weighted MRI. Journal of neuroscience, 31(32), 11597-11616.
6. Glasser, M. F., Smith, S. M., Marcus, D. S., Andersson, J. L., Auerbach, E. J., Behrens, T. E., ... & Van Essen, D. C. (2016). The human connectome project's neuroimaging approach. Nature neuroscience, 19(9), 1175-1187.
7. Huntenburg, J. M., Bazin, P. L., Goulas, A., Tardif, C. L., Villringer, A., & Margulies, D. S. (2017). A systematic relationship between functional connectivity and intracortical myelin in the human cerebral cortex. Cerebral cortex, 27(2), 981-997.
8. Mansour, S., Tian, Y., Yeo, B. T., Cropley, V., & Zalesky, A. (2021). High-resolution connectomic fingerprints: Mapping neural identity and behavior. NeuroImage, 229, 117695.
9. Margulies, D. S., Ghosh, S. S., Goulas, A., Falkiewicz, M., Huntenburg, J. M., Langs, G., ... & Smallwood, J. (2016). Situating the default-mode network along a principal gradient of macroscale cortical organization. Proceedings of the National Academy of Sciences, 113(44), 12574-12579.
10. Markello, R. D., Hansen, J. Y., Liu, Z. Q., Bazinet, V., Shafiei, G., Suárez, L. E., ... & Misic, B. (2022). Neuromaps: structural and functional interpretation of brain maps. Nature methods, 19(11), 1472-1479.
11. Raut, R. V., Snyder, A. Z., & Raichle, M. E. (2020). Hierarchical dynamics as a macroscopic organizing principle of the human brain. Proceedings of the National Academy of Sciences, 117(34), 20890-20897.


subj. feats.



atlas



spatial similarity




myelin maps


functional gradient (1)

similarity to grad. 1

Spatial similarity when comparing feature maps (myelin) to a common reference. With a map per subject, patterns can be interrelated (below). Sorting by overall similarity reveals a core-periphery structure[3], where maps with the lowest similarity to grad. 1 are dissimilar to others.




Sort by overall similarity to atlas




Difference in spatial similarity between the top and bottom correlated subjects (n=10). Bottom subjects load more onto the PCC whereas top subjects are more similar overall, and esp. in unimodal areas.

2

group-average spatial similarity: myelin to functional gradient 1





Spatial similarity can highlight locations of differential similarity. E.g., in orange we see the retrosplenial cortex, which has high myelin content and yet is considered transmodal according to gradient 1. This discordance is shown as dark blue.




Here we compute similarity reliability. We can compare maps to see where within-subject similarity is greatest, using *time series autocorrelation* half-width at half-maximum[11] as the exemplar feature.

Subject specific *day 1* to *day 2* similarity maps thresholded at top 20% of values






Group-average spatial similarity map showing where within-subject similarity is in the top 20% most often




3




This scatter shows how the locations of the feature extrema are where the between-day similarity is often occurring.

- For a given feature map, extrema values are generally co-located
- To find individual-specific information in similarity maps, an approach that scales for location-specific similarity is needed.

3+




Repeat this z-scoring operation for each of the ~20k vertices to make map per subject.



Subject-level discriminability maps

Group-averaged discriminability maps



Note the different spatial topography of within-subject similarity map (previous panel) compared to the discriminability map (right). The discriminability map highlights areas where subjects have self-similarity that exceeds blue distribution (above)[4,8].

Discussion

- There are many (infinite!) similarity patterns that can result in a correlation of a specific magnitude (e.g., r=0.46)
- Rather than measuring similarity with one r value, we can use the correlation unwrapping approach to see how similarity loads across the cortex. This information can be used for:
  - Exploratory data analysis, to better understand where maps relate highly or to spot potential biases. In our examples we can see how comparing PET feature maps might be influenced by SnR in insular regions, or how the retrosplenial cortex is an area of notable dissimilarity when comparing myelin and the primary functional gradient.
  - Quantitatively describing different similarity patterns, whether via comparing patterns or by deriving a feature of the spatial similarity maps that co-varies with a behavior or phenotype. Comparing myelin-to-grad1 similarities across subjects shows that low-correlation subjects have mapped patterns that are more dissimilar to all other patterns.
- Future work will look to relate this approach to other neuroimaging methods, like VBM or spatial registration, to see how spatial similarity can be used to complement other sorts of mapped changes (especially for the case of warp fields).
- Additional future work will seek to relate spatial similarity patterns, as well as discriminability maps, to behavioral and phenotypic information. We seek to use similarity to narrow down the patches where this information resides, which could potentially boost statistical power